

B-chromosome evolution

Juan Pedro M. Camacho^{1*}, Timothy F. Sharbel^{2†} and Leo W. Beukeboom³

¹*Departamento de Genética, Universidad de Granada, 18071 Granada, Spain*

²*Arbeitsgruppe Michiels, Max-Planck-Institut für Verhaltensphysiologie, PO Box 1564, D-82319 Seewiesen, Germany*

³*Instituut voor Evolutionaire en Ecologische Wetenschappen, Rijksuniversiteit Leiden, PO Box 9516, NL-2300 RA Leiden, The Netherlands*

B chromosomes are extra chromosomes to the standard complement that occur in many organisms. They can originate in a number of ways including derivation from autosomes and sex chromosomes in intra- and interspecies crosses. Their subsequent molecular evolution resembles that of univalent sex chromosomes, which involves gene silencing, heterochromatinization and the accumulation of repetitive DNA and transposons. B-chromosome frequencies in populations result from a balance between their transmission rates and their effects on host fitness. Their long-term evolution is considered to be the outcome of selection on the host genome to eliminate B chromosomes or suppress their effects and on the B chromosome's ability to escape through the generation of new variants. Because B chromosomes interact with the standard chromosomes, they can play an important role in genome evolution and may be useful for studying molecular evolutionary processes.

Keywords: B chromosomes; transposons; evolution; heterochromatin; repetitive DNA; Muller's ratchet

1. INTRODUCTION

Eukaryotic genomes are composed not only of genes found in normal chromosomes (A chromosomes) but also of myriads of selfish genetic elements which do not obey Mendelian laws of inheritance. Notable among these elements are the transposons, segregation distorters, various cytoplasmic factors and B chromosomes. Of these, the latter were really the first selfish genetic elements to be described (Wilson 1907), although their parasitic nature (Östergren 1945; Nur 1966, 1977) and selfishness (Jones 1985; Werren *et al.* 1987) were recognized many years following their initial descriptions.

The B chromosomes, also referred to as supernumerary or accessory chromosomes, are 'additional dispensable chromosomes that are present in some individuals from some populations in some species, which have probably arisen from the A chromosomes but follow their own evolutionary pathway' (J. P. M. Camacho & J. S. Parker, in Beukeboom 1994a). In addition, their irregular mitotic and meiotic behaviour allows them to accumulate selfishly in the germline, enabling non-Mendelian inheritance with transmission rates exceeding those of normal chromosomes (0.5). They have been found in all major groups of animals and plants.

B chromosomes have traditionally attracted much interest and various aspects of their biology have been reviewed several times (for an overview, see Jones 1995), of which the most recent comprehensive treatise is the monograph by Jones & Rees (1982). In this review, we discuss current insights into B-chromosome evolution and

point out new developments and directions in B-chromosome research that have occurred during the last two decades. A large number of recent studies have revealed new features of B chromosomes and some have shed light on previously unanswered questions. Among them are descriptions of previously unknown mechanisms of B-chromosome inheritance, an extreme example being the paternal sex-ratio (PSR) chromosome of the wasp *Nasonia*, which accumulates through its ability to destroy paternal chromosomes (Werren 1991). Long-term studies on specific B-chromosome systems (e.g. the grasshopper *Eyprepocnemis plorans*; Camacho *et al.* 1997a,b) have provided evidence for ongoing interactions between B-chromosome morphs and the standard genome at the level of local populations. Indeed, B-chromosome evolution may be viewed as the outcome of continuous conflict between parts of the genome with different interests, i.e. B-chromosome influences may shift back and forth from parasitic to neutral and possibly beneficial effects. Undoubtedly, the most important progress comes from the development and application of new molecular techniques. Data obtained from molecular analyses of several B chromosomes are now available to shed light upon questions of their origin and subsequent chromosomal evolution and sophisticated *in situ* hybridization techniques have additionally contributed to a better understanding of B-chromosome DNA composition and organization.

2. ORIGIN

In recent years, the types of DNA sequences residing on B chromosomes have been analysed extensively in some organisms. The first analyses in the 1970s and 1980s demonstrated that B chromosomes contained DNA that

* Author for correspondence (jpmcamac@ugr.es).

† Present address: Max-Planck-Institut für Chemische Ökologie, Tatzendpromenade 1a, 07745 Jena, Germany.

Table 1. *Classification of the origins of B chromosomes*

(At the highest classification level, B chromosomes are distinguished based upon their origination from the genome of their actual host species (intraspecies) or from the genome of another species (interspecies or hybrid origin). The second level of classification differentiates between origin from standard A chromosomes (autosomes) and sex chromosomes. For organisms with heteromorphic sex chromosomes, a further distinction can be made between origin from the homogametic or heterogametic sex chromosome. Evidence: 1, similar banding patterns of B chromosomes found among standard complement (including sex chromosomes); 2, (repetitive) DNA sequences of B chromosomes shared with standard complement (including sex chromosomes); 3, (repetitive) DNA sequences of B chromosomes not shared with standard complement, but present in related species; 4, direct observation of origination of chromosomal fragments in interspecific crosses; and 5, phylogenetic analysis of transposable elements shared by B chromosomes and standard complement (intra- and interspecies). See Beukeboom (1994a) and Hackstein *et al.* (1996) for details on the techniques used.)

classification	species	evidence	reference
intraspecific autosomal	<i>C. capillaris</i>	2	Jamilena <i>et al.</i> (1994, 1995)
	<i>S. cereale</i>		Jones & Flavell (1983), McIntyre <i>et al.</i> (1990), Sandery <i>et al.</i> (1990), Blunden <i>et al.</i> (1993), Cuadrado & Jouve (1994), Houben <i>et al.</i> (1996)
	<i>Z. mays</i>	2	Peacock <i>et al.</i> (1981), Viotti <i>et al.</i> (1985), Alfenito & Birchler (1993), Stark <i>et al.</i> (1996)
	<i>C. plumosus</i>	1	Keyl & Hägele (1971)
	<i>D. subsilvestris</i>	2	Gutknecht <i>et al.</i> (1995)
	<i>Petauroides volans</i>	2	McQuade <i>et al.</i> (1994)
	<i>Reithodontomys megalotis</i>	2	Peppers <i>et al.</i> (1997)
intraspecific sex chromosomal no heteromorphy	<i>E. plorans</i>	2	Leach <i>et al.</i> (1995), Houben <i>et al.</i> (1997a)
			López-León <i>et al.</i> (1994)
intraspecific sex chromosomal homogametic	no examples		
intraspecific sex chromosomal heterogametic	<i>Glossina</i> spp.	2	Amos & Dover (1981)
	<i>L. hochstetteri</i>	2	Sharbel <i>et al.</i> (1998)
interspecific autosomal	<i>Coix</i>	2,4	Sapre & Deshpande (1987)
	<i>N. vitripennis</i>	3,5	McAllister & Werren (1997)
	<i>P. formosa</i>	4	Schartl <i>et al.</i> (1995)
interspecific sex chromosomal no heteromorphy	no examples		
interspecific sex chromosomal homogametic	no examples		
interspecific sex chromosomal heterogametic	no examples		

was similar to that found on the A chromosomes (for a review, see Jones & Rees 1982). Research in the 1990s has involved the isolation, cloning and sequencing of numerous repetitive DNAs located on B chromosomes of various species. Some of these are specific to the B chromosomes while others are shared with the A chromosomes (reviewed in Beukeboom 1994a; Hackstein *et al.* 1996).

The traditional view, which is still widely accepted, is that B chromosomes are derived from the A chromosomes (Jones & Rees 1982). From this perspective, we could consider the origin of the B chromosome as a simple by-product of the evolution of the standard karyotype. For example, a B chromosome could derive itself from polysomic A chromosomes, from centric fragments resulting from A-chromosome fusions or from amplification of the paracentromeric region of a fragmented A chromosome. The first clear evidence in favour of the latter hypothesis was obtained by Keyl & Hägele (1971), who demonstrated that the polytene band pattern in the B chromosome of *Chironomus plumosus* was similar to that found near the centromere of chromosome IV.

Recent cytological and molecular studies support the notion that most B chromosomes seem to be derived from the autosomal complement of their current host species, but these studies have also demonstrated other modes of B-chromosome origin (table 1). Intraspecific origin from

A chromosomes is the most likely scenario for eight B chromosomes investigated, given the identification of similar repeat DNA sequences on them. For example, all repetitive DNA sequences isolated by micro-dissection from the B chromosome in *Crepis capillaris* are also present in the A chromosomes, although it has not been possible to identify from which autosome the B chromosome originated (Jamilena *et al.* 1994, 1995).

Sex chromosomes have previously been proposed as ancestors of B chromosomes since they may be more easily tolerated in the polysomic state (Hewitt 1973a). An example of a sex-chromosome-derived B chromosome is the B₂ chromosome of the grasshopper *E. plorans*, where the arrangement of two DNA sequences (a 180 bp tandem repeat and ribosomal DNA) with respect to the centromere coincide specifically with that of the X chromosome (López-León *et al.* 1994). This suggests that the B chromosome of *E. plorans* has been derived from the paracentromeric region of the X chromosome, with subsequent amplification of the two types of sequences contained there. Another example is the B chromosome of the New Zealand frog *Leiopelma hochstetteri*, which appears to be derived from a univalent (heteromorphic) W sex chromosome based on DNA sequence comparisons (Sharbel *et al.* 1998) and morphological similarities with the univalent W chromosome (Green *et al.* 1993).

The idea that B chromosomes in one species could have originated from the A chromosomes of a closely related species, which was originally proposed by Battaglia (1964), has recently gathered strength (Sapre & Deshpande 1987; McVean 1995; Schartl *et al.* 1995). Sapre & Deshpande (1987) demonstrated the spontaneous origin of B chromosomes in interspecific crosses between *Coix aquaticus* and *C. gigantea*. Possible evidence of the origin of B chromosomes by interspecific hybridization was recently found in the gynogenetic fish *Poecilia formosa*, a hybrid species between *P. mexicana* and *P. latipinna*. This unisexual species requires sperm of a sexual parental species to initiate egg development, but paternal chromosomes are eliminated from the developing zygote (Dawley 1989). Laboratory crosses between individuals of *P. formosa* and males of a black strain, both lacking B chromosomes, produced some black-pigmented offspring (frequency = 0.001), most likely the result of paternal pigmentation genes located on B chromosomes which appeared in the offspring because of incomplete elimination of paternal A chromosomes (Schartl *et al.* 1995).

The detection of DNA sequences that are restricted to the B chromosome of one species, but found on the A chromosomes of a closely related species would imply an interspecies origin. Although such a situation was previously reported for a B chromosome in *Brachycome dichromosomatica* (John *et al.* 1991), the repeat was later discovered at very low copy number in the standard genome (Leach *et al.* 1995). The best documented case of hybrid origin is that of the PSR B chromosome of the wasp *Nasonia*. McAllister & Werren (1997) used a phylogenetic analysis of DNA sequences of a retrotransposable element to show that the copies on the PSR were most similar to those of the species from the closely related genus *Trichomalopsis*. Hybridization studies with a linear piece of DNA from the PSR further supported a hybrid origin. Further phylogenetic analyses of sequences that are shared by B chromosomes of different species, such as those reported in the genus *Brachycome* (Houben *et al.* 1997a), may prove useful in elucidating the evolutionary history of B chromosomes.

Reproductive mechanisms that are based on chromosome elimination, as in the above-mentioned *Poecilia*, may be especially conducive to the origination of B chromosomes. The frequent occurrence of aneuploidy among sperm-dependent parthenogenetic (= gynogenetic) organisms (Beukeboom & Vrijenhoek 1998) supports this idea. Recently, Sharbel *et al.* (1997) described three different B-chromosome morphs in one population of the sperm-dependent parthenogenetic flatworm *Polycelis nigra*. Because the mechanism of sperm chromosome expulsion is imprecise (Benazzi Lentati 1970; Beukeboom *et al.* 1996), these B chromosomes may have originated from incompletely expelled autosomes. Similarly, the PSR B chromosome of *Nasonia* may have originated as a paternal fragmented chromosome following an escape from sperm chromosome destruction due to cytoplasmic incompatibility between *Nasonia vitripennis* and a species of *Trichomalopsis* (McAllister & Werren 1997). Although there are alternative explanations to cytoplasmic incompatibility, Ryan *et al.* (1985, 1987) showed that fragments of paternal chromosomes sometimes survive in cytoplasmic incompatible crosses.

3. MOLECULAR EVOLUTION

At the time of their origin, B chromosomes would be expected to share sufficient sequence and structural homology with their progenitor chromosomes such that they could synapse and recombine. However, their independent evolution and differentiation through processes analogous to Muller's ratchet require genetic isolation from any such elements within the nucleus (Green 1990; Beukeboom 1994a). It follows that newly arisen B chromosomes must have some predisposition to undergoing the relatively rapid structural modification required to induce synapsis failure. Intraspecific B-chromosome origin, which is probably the prevalent mode compared with those associated with hybridization, therefore presents a conundrum. If a neo-B chromosome originates from another chromosome, what initially inhibits synapsis between the two related chromosomes and allows the B chromosome to begin its journey towards independent evolution? Although these initial processes of chromosome evolution remain largely unknown, some indications are provided by data on polysomy in grasshoppers (Peters 1981; Talavera *et al.* 1990). These extra chromosomes, which are restricted to the germline and not inherited, are generated *de novo* each generation from autosomes through non-disjunction. In addition, they are heteropycnotic and do not pair with the original A chromosome. This suggests the presence of some cellular mechanism which can cause rapid heterochromatinization of extra elements (*sensu* genomic imprinting; Thomas 1995) and this could constitute the basis for B-chromosome differentiation (see also Hewitt 1973a). In contrast, a chromosome fragment crossing a species boundary will likely be sufficiently different to inhibit ectopic pairing with its new chromosome complement and, thus, such an element would immediately be univalent and prone to evolve as a B chromosome.

Subsequent to synaptic isolation within their respective genomes, elements of both types of origin (i.e. intra- and interspecific) will follow similar paths of molecular evolution and be subject to the same processes which act upon non-pairing chromosomes (Charlesworth 1978). B chromosomes thus converge upon a characteristic degenerate morphology, a reflection of the processes acting upon them rather than their mode of origin (Green 1990). Encouragingly, we are gaining some insights into the molecular evolutionary processes that act upon chromosomes once they have become isolated from the rest of the genome, in particular from studies of sex chromosomes and we discuss how these processes may play a role in B-chromosome evolution.

(a) DNA repeat sequences

B chromosomes are typically composed of repeated DNA sequences which vary dynamically in terms of repeat type and copy number (Amos & Dover 1981; Matzke *et al.* 1990; Sandery *et al.* 1990; Eickbush *et al.* 1992; Zeyl & Green 1992; Wilkes *et al.* 1995; Franks *et al.* 1996), a result of unequal crossing over and reduced recombination (Charlesworth *et al.* 1986; Stephan 1987). Repeats may form a significant part of the B-chromosome genome, as has been shown with different repeat families on the PSR (Eickbush *et al.* 1992; McAllister & Werren

1997) and, in some cases, repeats may be the exclusive constituent, as with the pSsP216 repeat unit in the B chromosomes of *Drosophila subsilvestris* (Gutknecht *et al.* 1995). McAllister & Werren (1997) have additionally shown that certain repeat sequences isolated from PSR of *N. vitripennis* are also found in the genus *Trichomalopsis*, thus providing evidence that they may be associated with mobile genetic elements (see § 3(c)). The typically heterochromatic nature of B chromosomes, as revealed by chromosome C banding, similarly demonstrates the presence of repeat DNA, as constitutive heterochromatin is generally composed of satellite blocks (Bigot *et al.* 1990; Charlesworth *et al.* 1994). In several cases, B chromosomes contain much larger amounts of repetitive DNA when compared to the genome from which they originated, thus suggesting massive amplification of repeat motifs over a relatively short time-scale, e.g. within one generation following a hybrid cross. It has also been suggested that repeat family amplification may be a mechanism through which a chromosome fragment (i.e. a neo-B chromosome) may become stabilized and positively selected for within a nucleus (Reed *et al.* 1994; Leach *et al.* 1995).

Repeat sequences have been implicated in lower vertebrate (Nanda *et al.* 1990, 1992, 1993) and plant (Guttman & Charlesworth 1998) sex-chromosome evolution and, thus, their influence upon B chromosomes may be analogous to the mechanisms leading to the evolution of heteromorphic sex chromosomes. In poeciliid fish, early sex-chromosome differentiation appears to have been initiated by the accumulation of simple repeat sequences adjacent to coding regions for sex determination (Nanda *et al.* 1990, 1992, 1993). In the white campion (*Silene latifolia*), a Y-chromosome-linked gene (MROS3) having an active X-chromosome-linked homologue appears to have been degenerated and silenced by multiple insertion–deletion events in addition to the accumulation of mononucleotide repeats (Guttman & Charlesworth 1998). Since the expression of this gene is limited to developing male flowers, a single active X chromosome copy in XY-chromosome plants is viable, thus alleviating any selection pressure to maintain function in the Y-chromosome homologue (Guttman & Charlesworth 1998). Genes on B chromosomes (assuming they were derived from transcriptionally active autosomal regions) are similarly under little or no selection pressure for maintenance of molecular genetic activity and, thus, they could probably undergo analogous suppressive changes through time. However, such genes still remain to be found on B chromosomes.

Once in position, repeat sequences may behave as nuclear protein targeting signals (Gilson *et al.* 1986; Charlesworth *et al.* 1994; Mitas *et al.* 1995) which can be highly specific, as evidenced by a protein that binds to heterochromatic autosomes but not to heterochromatic B chromosomes in male mealybugs (Epstein *et al.* 1992). Protein association with such sequences has been suggested as a mechanism through which significant conformational change in chromatid structure is established and efficient pairing with homologous regions of a sister element is prevented, effectively isolating these regions from recombination (Nanda *et al.* 1993). Mammalian X-chromosome inactivation is similarly mediated by chromatin–protein

association, as evidenced by the histone protein variant mH2A, which binds to non-heterochromatic regions of the X chromosome and probably causes changes in chromatin structure to induce transcription silencing (Costanzi & Pehrson 1998).

Finally, the accumulation of GACA and GATA repeats has been associated with the qualitative differentiation of cytologically indistinguishable sex chromosomes in the fish *Poecilia velifera* and *Xiphophorus maculatus* (Nanda *et al.* 1993). Southern hybridization of repeat sequence probes to genomic blots of closely related species lacking discernible sex chromosomes has demonstrated that many different types of repeat sequences have independently accumulated on the sex chromosomes of this group (Nanda *et al.* 1993). Thus, as a general mechanism through which synapsis between undifferentiated sex chromosomes can be inhibited, the exact repeat motif involved may be less important than the actual accumulation of microsatellite DNA itself. The accumulation of repeat sequences with subsequent meiotic isolation through conformational change in DNA structure may thus be the initiators of early heteromorphic sex-chromosome differentiation (Nanda *et al.* 1990, 1993) and these may represent plausible mechanisms through which intraspecific neo-B chromosomes are able to differentiate rapidly from their homologue progenitors.

(b) *Ribosomal DNA*

One form of tandemly repeated DNA which has been frequently described from B chromosomes is rDNA (see Green 1990). These genes, which encode ribosomal RNAs and exist as clusters of repetitive units, are typically visualized as secondary constrictions (nucleolar organizer regions or NORs) on metaphase chromosomes (e.g. through silver staining). Interesting insights into both B-chromosome origin and evolution may be made from rDNA.

It has been suggested that NOR regions are prone to chromosome breakage and this may provide a mechanism through which B chromosomes can be generated. NOR regions typically exhibit different times of expression (see Dai *et al.* 1994; Lin *et al.* 1995) relative to other autosomal genes and species-specific differences in rDNA condensation have been proposed as having led to the formation of a neo-B chromosome in a somatic hybrid between *Solanum brevidens* and *S. tuberosum* (Mitchell McGrath & Helgeson 1998). This process appears to have also acted in the genus *Brachycome* to generate different rDNA-containing B chromosomes (Houben *et al.* 1997a).

Chromosome regions containing rDNA show dynamic variation in repeat numbers and this has been attributed to deletions, duplications and unequal sister homologue exchange (Garrido *et al.* 1994; Garrido-Ramos *et al.* 1995). Intrahomologue recombination has additionally led to biased excision of rDNA between the recombining units in *Neurospora* (Butler & Metzenberg 1989). Assuming intrahomologue recombination and excision to be ubiquitous processes acting on rDNA clusters, the extent of NOR contraction is clearly limited in terms of organismal viability in autosomal regions but would potentially be under little or no selection on B chromosomes. A B chromosome which originates as an autosomal fragment containing an NOR region may lose its rDNA through

intrachromosome recombination and this may partially explain how B chromosomes degenerate, as variation in the number of rDNA repeats may significantly influence chromosome size (Adam 1992; Pukkila & Skrzynia 1993). This scenario does not necessarily exclude the possibility that the presence of rDNA on B chromosomes may give some selective advantage to B-chromosome carriers (Beukeboom 1994a); it could be that any selective advantage of having an rDNA-containing B chromosome simply decreases through time as rDNA copy numbers are decreased. Such a mechanism leading to the overall loss of rDNA repeat units may explain what have been considered anomalous results in repeated studies of NORs in different B-chromosome systems (Jones *et al.* 1989; Wilkes *et al.* 1995).

(c) *Transposable elements*

It has been proposed that B chromosomes might accumulate DNA from various sources (Beukeboom 1994a) existing as amalgamations of transposable DNA. This has been suggested as a mechanism through which some of the variability in mammalian Y chromosomes has arisen, as random insertions of transposable DNA into different regions of the Y chromosome would result in elements differing with respect to DNA composition and structure (Marshall Graves 1995). Compelling evidence for early Y-chromosome structural modification and allele silencing resulting from transposable element insertion comes from the TRAM element of the neo-Y chromosome of *Drosophila miranda* (Steinemann & Steinemann 1997). Transposable elements also appear to be involved in ectopic recombination (Montgomery *et al.* 1991), providing a plausible pathway through which sequences may be transferred across different homologues.

Theoretically, transposons should accumulate in regions not subject to recombination (Zeyl & Bell 1996) and this is supported by the transposable elements TRIM and TRAM the copy number of which on the neo-Y chromosome of male *D. miranda* (which undergoes recombination) is comparable to that of the complete female genome (Steinemann & Steinemann 1991, 1992; Steinemann *et al.* 1993). Clear evidence of a B chromosome providing a safe haven for a mobile element comes from work on the retrotransposon NATE (NAsonia Transposable Element) which has been described from the PSR element of *N. vitripennis* (McAllister 1995; McAllister & Werren 1997). A retrotransposon has also been invoked in the transposition of chloroplast DNA into the repeat element Bd49 of the B chromosomes of *B. dichromosomatica* (Franks *et al.* 1996). Mobile element insertion may thus be responsible for the generation of structural variability in B chromosomes. This mode of differentiation should proceed in a stepwise manner, with a B chromosome arising through the duplication of a major element followed by transposable element insertion. A duplicated autosomal region found on a B chromosome could thus rapidly lose homology with its parental sequence, the overall result being suppressed recombination between them. Such a scenario may be a contributing factor to the difficulties in elucidating B-chromosome origins.

Finally, any active genes inherited from the original progenitor elements of the B chromosomes may become

silenced either by insertions of transposable elements within the gene or through disruptions in ordered chromatin structure, as has been shown with silenced larval cuticle protein (*Lcp*) genes on the neo-Y chromosome of *D. miranda* (Steinemann *et al.* 1993). B chromosomes which are transcriptionally active and have no apparent phenotypic effects (Green *et al.* 1993) could thus conceivably have had their transcripts nullified through transposon insertion.

(d) *Epigenetic changes in B chromosomes*

Stem-loop structures are good candidates for protein binding sites and have been associated with heterochromatin condensation in the hymenopterans *Diadromus pulchellus* and *Eupelmus mullleti* (Bigot *et al.* 1990). The highly heterochromatic nature of B chromosomes may therefore (in part) be attributed to the presence of such secondary DNA structures. In the PSR chromosome, small palindromic sequences are associated with exchanges between repeats, suggesting that they enhance recombination between repeat units (Reed *et al.* 1994). Support for the potential of B-chromosome DNA to form hairpins *in vivo* also comes from the micro-dissected B chromosomes of the frog *L. hochstetteri* (Sharbel *et al.* 1998).

At the structural level, the single-stranded loop component of a hairpin may be prone to nucleolytic degradation (Mitas *et al.* 1995), a process that has been connected with chromosome breakage (Chen *et al.* 1995). For example, a G/CTT consensus sequence for topoisomerase I is found in certain eukaryote loop structures where single-stranded DNA cleavage occurs. The formation of hairpin structures in B chromosomes may attract DNases (Vogel *et al.* 1990), and expose enough single-stranded DNA to induce single-stranded cleavage and chromosome breakage, predisposing them to chromosomal rearrangements.

Methylation is hypothesized to cause sex-chromosome inactivation (Holliday 1987) and may therefore play a role in B-chromosome evolution. The B chromosome repeat family Bd49 of the Australian daisy (*B. dichromosomatica*) is hypermethylated and, thus, transcriptionally inactive and this is supported by an absence of Bd49 transcripts in leaf RNA extractions (Leach *et al.* 1995). In addition to effects on transcription, non-Mendelian B-chromosome behaviour may also be influenced by methylation. Neves *et al.* (1992) showed that induced demethylation (or blocking of methylation) in rye (*Secale cereale*) causes B chromosomes to undergo mitotic non-disjunction, a known B-chromosome accumulation mechanism.

Finally, chromatin packaging probably influences transcriptional regulation and the relative acetylation of histone molecules causes gene silencing in several organisms (see Houben *et al.* 1997b; Costanzi & Pehrson 1998). Houben *et al.* (1997b) showed that the B chromosomes of *B. dichromosomatica* are underacetylated relative to the autosomes and that this, in conjunction with late replication of B-chromosome DNA, may cause B chromosomes to become genetically inert. Similarly, autosomal rDNA transcription in *Allium* during different stages of mitosis is blocked by chromosome condensation (González-Fernández *et al.* 1993). Genetic inactivity of B chromosomes may thus not only be considered in terms of non-coding or non-functional DNA, but also from the

perspective of the many protein–DNA complexes which can be physically affected by chromatin structure.

It is becoming increasingly evident that the apparent similarities between B and sex chromosomes are more than just coincidental and that the molecular evolution of B chromosomes may therefore be interpreted in the context of sex-chromosome evolution (see Appendix A). Initially, a process must be available which can isolate a newly formed progenitor B chromosome relatively rapidly, such that homologous (or homoeologous) pairing is prevented. Such processes may be found in those analogous mechanisms which act upon heteromorphic sex chromosomes which have been studied in more depth. Subsequent to their isolation within the nucleus, B chromosomes would be expected to degenerate, in both structure and DNA sequence composition. This will make the identification of their progenitors more difficult over time.

4. FREQUENCY

B chromosomes have been described in more than 1300 species of plants and almost 500 species of animals (for reviews, see Jones & Rees 1982; Jones & Puertas 1993; Jones 1995) and in various species of fungi (Mills & McCluskey 1990; Miao *et al.* 1991a,b; Tzeng *et al.* 1992; Geiser *et al.* 1996; Leclair *et al.* 1996). These chromosomes have been described predominantly from certain taxonomic groups, although the high frequency of B chromosomes in these taxa probably reflects the intensity and technical ease with which each group has been studied. It is not surprising, therefore, that B chromosomes have frequently been reported in the Gramineae, Liliaceae and Orthoptera, groups which satisfy both these conditions. In fact, the discovery of B chromosomes in fungi was possible only after the development of a pulse-field gel electrophoresis technique for karyotyping these organisms. Thus, it is likely that many more species, when analysed with sufficient intensity, will be found to possess B chromosomes.

B chromosomes can attain extremely high frequencies in natural populations, depending both on the degree to which a particular species can tolerate these additional elements and on the strength of the B chromosomes' accumulation mechanism (if there is one). A stable frequency of B chromosomes is often found for several years in the same population, which has prompted authors in the past to conclude that polymorphism is in a state of equilibrium and that the frequency is a result of the action of two opposing forces—the accumulation of the B chromosome (which tends to increase the B-chromosome frequency) and the harmful effects on the fitness of the individuals carrying the B chromosome (which tend to decrease the frequency). However, as will be described later, B-chromosome polymorphism may be best interpreted as a dynamic system in which the frequency continually shifts due to an arms race between the A and B chromosomes.

In addition, interpopulational differences in B chromosome frequency depend on selective factors (i.e. the ecological tolerance of B chromosome carriers in terms of the permissiveness of the environmental conditions for a particular population), historical factors (i.e. the number of generations since B-chromosome origin), transmission factors (related to the differences between populations in

the accumulation intensity of the B chromosome) and random factors (i.e. the action of genetic drift in populations of finite size). The four types of factor probably act simultaneously, making it difficult to evaluate the relative importance of any single one, even under intense analysis. Some insight into the relative importance of these factors may be obtained from the distribution of B-chromosome counts among individuals sampled from one population analogous to tests on transposable elements in *Drosophila* (Charlesworth & Lapid 1989; Charlesworth *et al.* 1992a,b).

The maximum number of B chromosomes that a species is capable of tolerating, measured by the maximum number of B chromosomes found in adult individuals, varies broadly, although it ultimately depends on the relative intensities of the above-mentioned factors. Corn plants have been found with 34 B chromosomes (involving a 155% increase in nuclear DNA content; see Jones & Rees 1982), a situation which is probably tolerable to the plant because of its domestication. In wild plants, such as *Lolium perenne* (Jones & Rees 1982) and *B. dichromosomatica* (Carter 1978), individuals have not been found with more than three B chromosomes, although in *Allium schoenoprasum* plants have been reported carrying up to 20 B chromosomes (Bougourd *et al.* 1995). In the grasshopper *E. plorans* (Camacho *et al.* 1997b) and the flatworm *P. nigra* (Beukeboom *et al.* 1996), it is also rare to find individuals with more than three B chromosomes in natural populations, while individuals of the endemic New Zealand frog *L. hochstetteri* can have up to 15 mitotically stable B chromosomes (Green *et al.* 1993).

5. EFFECTS

Most B chromosomes are heterochromatic, promoting the general idea that these elements are genetically inert. Analyses of general transcriptional activity using tritiated uridine have supported this idea (Fox *et al.* 1974; Ishak *et al.* 1991). Nevertheless, some B chromosomes show transcriptional activity, as has been shown in the plumose state in the frog *L. hochstetteri* (Green 1988) or in the polytene state of the mosquito *Simulium juxtacrenobium* (Brockhouse *et al.* 1989). In addition, many B chromosomes have been found to carry ribosomal genes (for reviews, see Green 1990; Beukeboom 1994a; Jones 1995), although they are for the most part inactive (Donald *et al.* 1997). Some effects of the B chromosomes appear to be attributable directly to the products of their genes, as is the case with genes controlling resistance to rust in the B chromosomes of *Avena sativa* (Dherawattana & Sadanaga 1973), and the genes conferring resistance to antibiotics in the B chromosomes of the fungus *Nectria haematococca*, thereby favouring its pathogenicity (Miao *et al.* 1991a,b). These examples indicate that not all B chromosomes are genetically inactive. However, much more information is needed to support the generally accepted opinion that most B chromosomes lack major genes.

There is ample evidence that B chromosomes can affect a multitude of cellular and physiological processes in both plants and animals. The effects are rarely manifest in the external phenotype, although the B chromosomes of *Haplopappus gracilis* influence the colour of the achenes (Jackson & Newmark 1960) and, in corn, plants with B

chromosomes develop striped leaves (Staub 1987). More frequently, B chromosomes affect processes or characters associated with vigour, fertility and fecundity. Jones & Rees (1982) summarized a broad range of mostly detrimental effects from the B chromosomes in many species of plants and animals. These negative influences on host fitness pointed to the parasitic nature of B chromosomes. Nevertheless, some B chromosomes, when present in low numbers, have beneficial effects upon their carriers and, thus, may have a different biological significance (discussed in §7(a)). For example, the B chromosomes of various species of plants are associated with increased germination vigour or speed (see table 4.2 of Jones & Rees (1982)).

The influence of B chromosomes may stem either from their presence or from the activity of genes found on them. For example, the B chromosomes of the plant *Scilla autumnalis* (Ruiz-Rejón *et al.* 1980; Oliver *et al.* 1982) and *A. schoenoprasum* (Plowman & Bougourd 1994) alter the expression of A-chromosome genes for an esterase and endosperm protein, respectively. The presence of B chromosomes can also influence the expression of NORs on the A chromosomes, as is the case for the grasshopper *E. plorans* (Cabrerero *et al.* 1987). As mentioned above, many B chromosomes contain ribosomal genes, the activity of which could give the cell higher levels of translation (but see Donald *et al.* 1997). It would therefore be informative to study the possible effect that B chromosomes possessing active NORs might have on growth rates.

It should be emphasized that B-chromosome effects depend on the environmental conditions acting upon a population and can be characterized by both spatial and temporal variation. It is therefore risky for the effects detected in one population to be extrapolated over the entire distribution range of that species. Each case should be analysed thoroughly in many populations and the effects should be studied under the most natural conditions possible.

6. TRANSMISSION

Given that B chromosomes do not always occur in pairs and segregate to opposite poles during meiosis (the behaviour that stabilizes chromosome number in A chromosomes), they do not conform to a Mendelian system. Non-B univalent chromosomes would be expected to have meiotic transmission rates of 0.5, but this is typically lower as they are unstable in meiosis and/or mitosis. Many B chromosomes register transmission rates clearly greater than 0.5, i.e. they show accumulation, the most important property of parasitic B chromosomes. Accumulation can take place before, during or after meiosis; Jones (1991) exhaustively reviewed the principal cytological mechanisms that cause this accumulation. Regarding a pre-meiotic mechanism, it suffices to mention B-chromosome accumulation in the locust *Locusta migratoria* derived from their mitotic instability and the preferential destiny of cells with a high number of B chromosomes to become spermatogonia (Nur 1969). Meiotic accumulation has been described from female meiosis in various species of plant and animal and is based on the inherent asymmetry in the production of only one ovule from each oocyte; the B chromosome

migrates preferentially to the secondary oocyte instead of to the first polar body. In the insect *Pseudococcus affinis*, the B chromosomes accumulate during male meiosis by escaping the heterochromatinization and elimination of a chromosomal set characteristic of spermatogenesis (Nur 1962). Post-meiotic accumulation is frequent in plants, where the formation of pollen grains involves two post-meiotic mitotic divisions that give rise to the generative and vegetative nuclei; the non-disjunction of the B chromosome in this mitosis and the preferential migration of the two B chromatids to the generative nucleus are responsible for B-chromosome accumulation.

There is even a case of ameiotic accumulation of a B chromosome in the parasitoid wasp *N. vitripennis* (Werren 1991), where the B chromosome (PSR) present in the spermatozoa causes the condensation and loss of the paternal chromosomes accompanying it, transforming the diploid (female) zygote to a haploid male carrying the B chromosome. Through this mechanism the B-chromosome's transmission rate approaches one and, because it reduces the fitness of its host to zero, this B chromosome is considered one of the most parasitic of all known genetic elements.

In two species, rye (for references, see Jones & Rees 1982) and *L. migratoria* (Pardo *et al.* 1994), B chromosomes accumulate through both sexes, whereas in the grasshopper *Myrmeleotettix maculatus* B chromosomes show drive through females but drag through males (Hewitt 1973*a,b,c*). However, not all B chromosomes show accumulation, as in the plants *Poa alpina* (Håkansson 1954), *P. trivialis* (Bosemark 1957), *Centaurea scabiosa* (Fröst 1958), *Ranunculus acris* (Fröst 1969), *A. schoenoprasum* (Bougourd & Parker 1979) and *Guizotia scabra* (Hiremath & Murthy 1986). In animals, the most notable case is that of the grasshopper *E. plorans*, in which the three most frequent types of B chromosomes lack accumulation mechanisms (López-León *et al.* 1992*a*). These examples suggest the existence of other models of B-chromosome evolution that differ from the parasitic one, as we shall discuss in §7.

7. DYNAMICS

In general, B chromosomes could be considered genome symbionts the population dynamics of which depend on two important properties, i.e. their effects on genome fitness and their transmission ratio. Several outcomes are theoretically possible from the interaction of these two properties (table 2). It is clear that, subsequent to their origin, B chromosomes require accumulation mechanisms, otherwise their proliferation may only be explained in terms of beneficial effects on carriers. These are the only ways in which these chromosomes can increase in frequency and establish a polymorphism in a natural population (categories 1, 4 and 7–9). A newly risen B chromosome falling into category 5 would fail to establish a polymorphism, as failure to synapse and irregular meiotic behaviour would preclude its ability to become fixed by genetic drift. As opposed to the A chromosomes and the genes they contain, which normally follow the laws of Mendelian inheritance, the B chromosome is destined to extinction through random forces. Therefore, a near-neutral B chromosome (category 5) is

Table 2. Possible outcomes of the effect on host genome fitness and B-chromosome transmission rate on the establishment of a B chromosome

(Δw_G , change in host genome fitness due to B chromosome presence. $k_B - 0.5$ indicates the difference between the B-chromosome transmission ratio (k_B) and the Mendelian one (0.5); therefore + indicates B chromosome accumulation and - indicates B chromosome elimination.)

category	Δw_G	$k_B - 0.5$	net result	evolutionary significance
1	-	+	B polymorphism	parasitism
2	-	0	B disappears	—
3	-	-	B disappears	—
4	0	+	B polymorphism ^a	attenuated parasitism
5	0	0	B disappears	near-neutral B chromosome
6	0	-	B disappears	—
7	+	+	B polymorphism ^a	mutualism
8	+	0	B polymorphism ^a	mutualism
9	+	-	B polymorphism	mutualism

^a Infinite accumulation is prevented because individuals with high numbers of B chromosomes have reduced fitness.

presumably derived from an attenuated parasitic B chromosome (category 4) that has lost drive or from a mutualistic B chromosome (e.g. category 8) that is no longer beneficial for genome fitness. If the new B chromosome were harmful to the carriers, it would only be able to persist if its propensity to accumulate outweighed any negative effects upon its carriers. This may explain the origin of most of the parasitic B chromosomes known today from a multitude of species (category 1).

(a) *Equilibrium models*

The two most widely accepted models of B-chromosome evolution, the heterotic model (White 1973) and the parasitic (Östergren 1945; Nur 1966, 1977) or selfish model (Jones 1985; Shaw & Hewitt 1990), assume that the frequencies of B chromosomes are in equilibrium in current populations and are used to contrast the antagonistic forces responsible for the equilibrium. The heterotic model assumes a balance between the positive fitness effects of B chromosomes (which show no accumulation) when they occur in low numbers and their negative effects when they occur in high numbers. Typically, it has been applied to category 8, but could equally fit categories 7 and 9 (table 2). The only known B chromosome which has a strong likelihood of being heterotic is that of the chive *A. schoenoprasum*. While this B chromosome does not show accumulation, it has been demonstrated that plants with B chromosomes survive better in natural habitats than those without B chromosomes (i.e. in terms of the development from seed to seedling; Holmes & Bougourd 1989) due to the fact that the B chromosomes boost the germination rate under drought conditions (Plowman & Bougourd 1994).

For the parasitic-selfish model, the equilibrium is the result of B-chromosome accumulation (which increases its frequency) and, typically, of its detrimental effects on the fitness of B-chromosome carriers (which reduce the

frequency of the B chromosome). The great majority of B-chromosome systems that have been analysed in detail fall into category 1 (table 2) and are thus compatible with the parasitic model (see Nur 1977; Jones 1985, 1995; Nur & Brett 1985, 1987, 1988; Ruiz-Rejón *et al.* 1987; Shaw & Hewitt 1990).

In most of these studies, B-chromosome frequencies are minimized through their increasing negative effects on host fitness when they increase in number, but other selective pressures may also play a role. A good example is the PSR chromosome of the parasitoid wasp *N. vitripennis*, in which population structure and fertilization proportion affect the spread of the PSR (Beukeboom & Werren 1992; Werren & Beukeboom 1993). The PSR has a transmission rate to sperm of nearly one but causes destruction of the paternal chromosomes, except for itself, shortly after egg fertilization. Owing to haplodiploidy, this results in the conversion of diploid (female) eggs to haploid (male) eggs that carry the PSR. *Nasonia vitripennis* parasitizes fly pupae that occur in temporary patches (e.g. at carcasses) resulting in a demic population structure where flightless males mate locally with their emerging sisters. A theoretical analysis showed that the PSR equilibrium frequency is strongly affected by deme size (the number of founding females) and the fertilization proportion. Population experiments under laboratory conditions confirmed most of the theoretical predictions, i.e. it led to loss of the PSR from populations consisting of small deme sizes and when the fertilization proportion was low. Although these laboratory results have not been repeated under natural conditions, Beukeboom (1994b) showed that the PSR causes such minor effects on various traits related to carrier fitness that population structure and fertilization proportion play the major role in determining the frequencies of the B chromosome in natural populations.

(b) *Tolerance to B chromosomes*

It has been a parasitological dogma that a well-adapted parasite should not damage its host, as debilitation and death of the host can cause the death of the resident parasites (Hoepflich 1977; Alexander 1981). However, theoretical (Anderson & May 1982; May & Anderson 1983) and comparative analyses (Ewald 1987) have suggested that this is not necessarily so. For instance, the evolution of parasite virulence (the effect of a parasite on host fitness) may be strongly influenced by the parasite's mode of transmission (Anderson & May 1982; Lipsitch *et al.* 1995), i.e. parasites transmitted horizontally should be more virulent than those transmitted vertically, because the latter have their fitness linked to the fitness of their host and, therefore, harming of the host will reduce parasite fitness. In contrast, horizontally transmitted ones can be more virulent because they may contagiously infect other individuals. Mathematical models predict that parasites that are only vertically transmitted should evolve towards less virulence (Lipsitch *et al.* 1995) and several comparative (Ewald & Schubert 1989; Herre 1993; Clayton & Tompkins 1994) and experimental (Bull *et al.* 1991) studies have shown that the degree of vertical transmission in nature is positively correlated with benignity.

B chromosomes are exclusively vertically transmitted parasites and, hence, fit the expectation of the evolution towards attenuated parasitism, e.g. a change from category 1 to 4 in table 2. This may result from the appearance of less virulent B-chromosome types and/or the evolution of more tolerant host genotypes. As Shaw (1984, p. 93) pointed out, 'alleles on the A chromosome set that reduce the selection operating against animals carrying B's will be selected, as will B-chromosomes that are less damaging to their carrier'. Although many detrimental effects of B chromosomes have been reported (Jones & Rees 1982), it should be borne in mind that the evolution of B-chromosome tolerance depends on the existence of appropriate genetic variation and a high increase in B-chromosome frequency, because selection for B-chromosome tolerance can only take place in B-chromosome-carrying individuals. It is thus conceivable that B-chromosome tolerance has not evolved in all known systems. The inability to detect significant effects of B chromosomes on carrier fitness would be consistent with the evolution of B-chromosome tolerance in a natural population. For instance, the locust *L. migratoria* (Castro *et al.* 1998) harbours attenuated parasitic B chromosomes that do not produce apparent deleterious effects on B-chromosome carriers (category 4). Likewise, the grasshopper *E. plorans* (López-León *et al.* 1992a,b; Camacho *et al.* 1997a,b; Martín-Alganza *et al.* 1997) possesses B chromosomes that were originally parasitic but whose drive has subsequently been neutralized by the host genome. These B chromosomes have no apparent effects on carrier fitness. Interestingly, a new parasitic B-chromosome variant (B₂₄) that has recently replaced the neutralized B-chromosome version (B₂) significantly reduced egg fertility (Zurita *et al.* 1998). This suggests that newly arisen parasitic B-chromosome variants are more harmful than older B-chromosome versions. These conclusions are preliminary since the evolution of tolerance to B chromosomes has not received much consideration in the past, mainly because of the difficulty in detecting slight effects. Nevertheless, there are a large number of B-chromosome systems where no significant B-chromosome effects have been detected.

(c) *Suppression of drive*

An absence of accumulation does not necessarily indicate that a B chromosome is heterotic. Parasitic B chromosomes impose a genetic load upon carrier populations and, thus, favour the evolution of any gene variants on the A chromosomes which would tend to reduce this load, either by eliminating B-chromosome accumulation (= B-chromosome resistance genes) or by buffering any detrimental effects (the evolution of B-chromosome tolerance genes; see §7(b)). The presence of some type of A-chromosome genetic control over B-chromosome accumulation has been demonstrated in *S. cereale* (Müntzing 1954; Romera *et al.* 1991; Jiménez *et al.* 1995), *Festuca pratensis* (Bosemark 1954), *Zea mays* (Carlson 1969; Rosato *et al.* 1996), *Hypochoeris maculata* (Parker *et al.* 1982), *M. maculatus* (Shaw & Hewitt 1985; Shaw *et al.* 1985), *Pseudococcus affinis* (Nur & Brett 1985, 1987, 1988), *Aegilops speltoides* (Cebriá *et al.* 1994) and *E. plorans* (Herrera *et al.* 1996). Such evidence has been extrapolated mostly from variation in transmission rates between individuals, from

the success of artificial selection in obtaining lines of high and low transmission rates and from the different results obtained through intra- and interpopulational crosses. Recently, evidence has been provided that B chromosomes in rye possess genes controlling their own transmission (Puertas *et al.* 1998).

(d) *A non-equilibrium model of long-term evolution*

Parasitic B chromosomes that have lost their accumulation mechanisms are doomed to disappear, unless they become heterotic or recover accumulation at some point during the long process towards random extinction, thereby transforming them into a new type of parasitic B chromosome. According to the magnitude of negative effects exerted by the B chromosomes at the time of losing accumulation, they would disappear rapidly (large effects), slowly (small effects) or very slowly (imperceptible effects) from the population. In this last case, we can consider B chromosomes to be near neutral (category 5), a type of B chromosome that is not found in equilibrium, but that, as we shall see below, constitutes a transitory stage towards disappearance or towards regeneration of the polymorphism.

The only proof of the existence of near-neutral B chromosomes, that is those that have lost accumulation and produced insignificant effects on the fitness of the carriers, is presently provided by the B chromosome of the grasshopper *E. plorans* (see Appendix B). While the transmission ratio of B chromosomes in *E. plorans* is usually close to 0.5 in most individuals (López-León *et al.* 1992a), these ratios can vary greatly between individuals in many species with parasitic B chromosomes (Bosemark 1954; Müntzing 1954; Parker *et al.* 1982; Nur & Brett 1985, 1987, 1988; Shaw & Hewitt 1985; Shaw *et al.* 1985; Romera *et al.* 1991; Cebriá *et al.* 1994; Jiménez *et al.* 1995). This could be due to the fact that the suppression of B-chromosome accumulation imposes negative pleiotropic effects of the genes involved, thereby impeding any marked increase in their frequency and preventing complete suppression of B-chromosome accumulation.

The B-chromosome system in *E. plorans* not only illustrates the presence of parasitic chromosomes neutralized by the A-chromosome genome, but goes further to provide evidence of one of the few evolutionary paths remaining for these B chromosomes (apart from disappearing, but this is of minor evolutionary interest); it involves the regeneration of the polymorphism through the appearance of a new parasitic B-chromosome variant that starts the cycle again (figure 1).

Overall, the polymorphism for the B chromosome of *E. plorans* has been regenerated on at least three occasions on the Iberian Peninsula (assuming that B₁ was the ancestral type, given that it is predominant in the majority of the populations analysed): (i) when B₁ was substituted by B₂ in the province of Granada and the eastern part of the province of Málaga, (ii) when B₁ was replaced by B₃ in the zone of Fuengirola (Málaga), and (iii) when B₂ was replaced by B₂₄ in Torrox. This polymorphism illustrates that B-chromosome polymorphisms must not be seen necessarily as a system in equilibrium, but rather as a dynamic succession of stages through which the same polymorphism can change from parasitism to near neutrality and then to parasitism again. Thus, the B




Figure 1. Arms race making up the long-term evolution of a parasitic B chromosome. The life cycle of the B chromosome begins with the parasitic stage in which the B chromosome possesses drive (which is the property facilitating its establishment in the population) and is harmful for the individuals carrying it. Neutralization consists of the suppression of B-chromosome drive by the evolution of appropriate genes in the A-chromosome genome, paralleled by the evolution of an A-chromosome genome more tolerant to B-chromosome effects and/or less harmful B-chromosome variants. The near-neutral B chromosome possesses no drive and causes minor effects on carriers, at least at low numbers. This leads it to a random walk towards stochastic loss (since B-chromosome fixation is impossible because of meiotic irregularity and harmfulness at high B-chromosome numbers) that may be long in large populations. However, if in the course of the random walk the B chromosome mutates to a new variant capable of driving, then the polymorphism regenerates and the cycle starts again.

chromosome's existence can be prolonged by dynamically resisting A-chromosome genome assaults which tend to force the B chromosome to disappear.

It is not possible at present to determine how many systems of B chromosomes might be similar to the system of *E. plorans*, but a thorough analysis of those that do not show accumulation or that show it in some populations but not in others will almost certainly show similar cases in the near future. The non-equilibrium model developed for the *E. plorans* B chromosomes illuminates the possible long-term evolution of not only parasitic B chromosomes but also other selfish genetic elements (Johnson 1997).

8. ROLE

The role of B chromosomes in the evolution of eukaryotic genomes appears at first to be somewhat superfluous. Given that their presence is not needed for survival or reproduction of the individual, these chromosomes appear at first to be simply 'genomic junk', a waste product of the eukaryotic genome. Nevertheless, given that B chromosomes cannot originate as junk (that is, as neutral or slightly harmful), but most probably begin by being selfish (because if they had no accumulation they could not increase in frequency), it is plausible that, of all the extra chromosomes produced over the evolution of genomes of most organisms, only the selfish ones can be transformed into B chromosomes.

We have already discussed the neutralization of a parasitic B chromosome, but one might ask whether a neutralized B chromosome could become heterotic. The idea that a parasitic B chromosome could become heterotic was proposed first by Kimura & Kayano (1961) and has also been defended by Ruíz-Rejón *et al.* (1987). The main problem arising from this possibility involves the fact that the genes of the B chromosomes are

generally inactive and can therefore accumulate a large number of mutations. One must then ask how the B chromosomes evolved a beneficial characteristic for the carrier, somehow resisting the effect of Muller's ratchet. There are several possible answers: (i) as we observed above, some genes of the B chromosomes are active, thus conserving some of their functional requirement, suggesting that under certain circumstances these genes may be advantageous to the genome, and (ii) B chromosomes can capture genes from the A-chromosome genome and, thus, become indispensable to the host. This is the case mentioned above for the gene present in the B chromosome of the fungus *N. haematococca*, which makes it resistant to pisatine, an antibiotic produced by its host plant, the pea (Miao *et al.* 1991a,b).

In the same light, it has recently been proposed that the Y chromosome of *Drosophila* may have evolved from a supernumerary chromosome with the characteristics of a B chromosome (see Hackstein *et al.* (1996) for details of the argument). This idea, which suggests one of the forms in which a B chromosome can ultimately integrate itself into the A-chromosome genome, makes sense only when the B chromosome provides more or less essential functions and achieves regularity in meiosis. For example, in the zebra finch *Taeniopygia guttata*, a supernumerary chromosome has recently been reported which is restricted to the germline such that all males and females carry one (Pigozzi & Solari 1998). The authors have proposed a mechanism by which one copy of the supernumerary is always present in all individuals, i.e. through complete elimination in male meiosis and total preferential segregation to the oocyte in females. If this hypothesis were correct, this could be a stabilized B chromosome transmitted through a single sex (the female), a case reminiscent of the *Nasonia* one. We could imagine the zebra finch situation as a possible solution for many selfish B chromosomes that show strong drive through the

female and are partly eliminated through the male, e.g. *M. maculatus* (see §6). Their absence from the somatic line avoids most of their harmful effects on carriers and their special transmission mechanism assures their regular presence in the germline. In addition, it is conceivable that B chromosomes can integrate themselves into the genome by translocation, a possibility first suggested by White (1973) and supported by the spontaneous translocations that have been recorded between the A and B chromosomes in *E. plorans* (Henriques-Gil *et al.* 1983; Cabrero *et al.* 1987). At first sight, the possibility that B chromosomes may become transformed into essential members of the A-chromosome genome (even forming part of some of the chromosomes of the A set) is a remote one. However, there are some recent data suggesting this possibility since a population analysis in the Brazilian wasp *Trypoxylon albirtarse* has shown that, in most populations in the Viçosa region, most males have one B chromosome and most females have two B chromosomes. Since males are haploid and females diploid, it seems that this B chromosome is close to stabilization in these populations (S. M. S. R. Araujo, personal communication). Perhaps these B chromosomes could represent only a fraction of the total heterochromatin of these species and their fate could depend on their possible role in the genome. The molecular analysis of this system is an interesting task for future research in the B-chromosome field.

9. PERSPECTIVES

We have discussed a variety of B-chromosome systems each with its particular transmission dynamics. Although most B chromosomes show accumulation of one form or another (Jones 1991), some exceptional B chromosomes transmit at nearly Mendelian rates and, therefore, cannot be categorized as selfish. This is the case, for example, with the B chromosomes of the chive *A. schoenoprasum* (Bougourd & Parker 1979) and of the grasshopper *E. plorans* (López-León *et al.* 1992a), studies which offer new perspectives on the evolution and biological meaning of these enigmatic chromosomes. We have seen the discovery of new B-chromosome systems with previously unknown mechanisms of (accumulation) transmission and we can expect more in the future.

Recent molecular studies of B chromosomes have revealed that they do not have a single mode of origin, but instead can arise in a variety of ways. Although the source of the B chromosomes can sometimes be traced with a degree of certainty (e.g. the autosomal complement, sex chromosomes or a closely related species), it has still been difficult to pinpoint the exact progenitor DNA region(s). The extent to which this will be possible is unclear at the moment. It will depend critically on intensive molecular study as well as our ability to determine the speed and nature of the molecular processes involved in chromosome evolution. This should also reveal why we cannot find any traces of silenced (relict) genes on existing B chromosomes (other than rDNA).

B chromosomes are being recognized as suitable systems for studying genome evolution. As single chromosomes that have been freed from the selection pressures that act on the maintenance of standard chromosomes,

they may prove useful in studying processes of molecular degeneration analogous to studies of heterogametic sex chromosomes. We expect that they will become important in understanding chromosomal evolution, including evolution of repetitive DNAs, gene silencing and transposable elements.

J.P.M.C. expresses his most sincere appreciation to all those who participated in the study of B chromosomes in *E. plorans* and particularly to J. Cabrero, M. D. López-León, M. C. Pardo and M. W. Shaw, without whose assistance we would have advanced very little in disentangling this enigma. T.F.S. and L.W.B. acknowledge the supportive and stimulating scientific environment of Arbeitsgruppe Michiels at the Max-Planck-Institut für Verhaltensphysiologie. In addition, we would like to thank J. Cabrero, M. D. López-León, U. Nur, J. L. Oliver, M. Ruiz-Rejón, J. L. Santos, M. W. Shaw and J. H. Werren for fruitful discussions on B-chromosome evolution and T.F.S. would in particular like to thank D. M. Green for introducing him to these wonderful genetic elements. This research would not have been possible without the financing awarded by the Dirección General de Investigación Científica y Técnica and the Junta de Andalucía to J.P.M.C. and stipends of the Deutsche Forschungsgemeinschaft to L.W.B. and T.F.S. and of the Royal Netherlands Academy of Arts and Sciences to L.W.B. We all are grateful to D. M. Green, B. F. McAllister and J. N. Timmis for their critical reading of the manuscript and two anonymous referees for useful comments.

APPENDIX A. REASONS FOR SIMILARITY BETWEEN SEX CHROMOSOMES AND B CHROMOSOMES

B chromosomes are often similar to sex chromosomes in terms of meiotic behaviour, size, morphology and heteropycnocity (Hewitt 1979; Amos & Dover 1981; Jones & Rees 1982; Green 1990), and this resemblance can be interpreted in a number of ways.

- (i) In the orthopteran, *Melanoplus femur-rubrum*, B-chromosome similarity to their X chromosomes has been explained in terms of chromosome inactivation, as the heteropycnocity of both elements is likely required for normal meiosis (Nur 1978). As such, the B and sex chromosomes of *M. femur-rubrum* converge upon a common morphotype due to the functional constraints of meiosis (i.e. to prevent pairing of B and X chromosomes with autosomal homologues; Dover & Riley 1972).
- (ii) Shared similarity between B and sex chromosomes may also imply real homology, as has been shown by Amos & Dover (1981), who demonstrated that the B chromosomes of the fly *Glossina* have arisen from a duplicated Y chromosome and have subsequently become differentiated from the Y chromosome through the accumulation of tandem repeat DNA or as in the frog *L. hochstetteri*, whose B chromosomes have been derived from the univalent W sex chromosome (Green *et al.* 1993; Sharbel *et al.* 1998).
- (iii) B chromosomes and univalent members of heteromorphic sex chromosomes may converge upon a typical degenerate morphotype due to similar molecular evolutionary processes acting upon them (Green 1990).

APPENDIX B. THE B-CHROMOSOME SYSTEM OF *E. PLORANS*

The B-chromosome polymorphism of *E. plorans* is extremely widely distributed over Mediterranean and southern Atlantic coastal regions of the Iberian Peninsula (Henriques-Gil *et al.* 1984), the north of Africa (Henriques-Gil 1984) and Italy (López-Fernández *et al.* 1992). Its most important characteristics are as follows.

- (i) The propensity to mutate, as exemplified by the high number of novel B-chromosome types described to date (more than 40 have been differentiated on the basis of size, morphology and C-banding; Henriques-Gil *et al.* 1984; Henriques-Gil & Arana 1990; López-León *et al.* 1993) and that new types of B chromosome can appear between the offspring of controlled crosses where none of the offspring carried this type of B chromosome (López-León *et al.* 1993). The most widely distributed B chromosome, called B₁, is considered the ancestral B chromosome from which the rest were derived (Henriques-Gil *et al.* 1984) by replacement processes (Henriques-Gil & Arana 1990).
- (ii) The three most frequent types, B₁, B₂ and B₅, lack accumulation (López-León *et al.* 1992a) and lack significant effects over several traits related to the fitness of the carriers (López-León *et al.* 1992b; Camacho *et al.* 1997a,b).
- (iii) B₂ is capable of accumulating in females crossed with males from populations having no B chromosomes, but not when the same females are crossed with males from the same population (Herrera *et al.* 1996), thus suggesting that the B chromosomes originally had an accumulation mechanism which was lost due to the evolution of drive suppressor genes in the A chromosomes, which, logically, are not found in the males of the population that lacks B chromosomes.

In the population of *E. plorans* captured in 1984 near Torrox (Málaga), Henriques-Gil & Arana (1990) verified the dominance of a B-chromosome type termed B₂₄ (with a mean number of B chromosomes of 0.344), which was different from the predominant type in adjacent populations (B₂). This new B chromosome was like B₂, but with a duplicated proximal band associated with a greater amount of repetitive 180 bp DNA than possessed by B₂, as well as a lesser amount of ribosomal DNA (Zurita *et al.* 1998). After also finding B₂, although at a very low frequency, Henriques-Gil & Arana (1990) proposed that B₂ was being replaced by B₂₄ in this population. In 1992, Zurita *et al.* (1998) captured specimens in this same location which showed a B-chromosome frequency of 0.975, enormously exceeding that of 1984 and made a series of controlled crosses which indicated that B₂₄ had a strong tendency to accumulate through females, their mean transmission ratio (0.696) being significantly higher than a Mendelian one. However, they did not find any traces of B₂ in the sample analysed from 1992 and it would thus appear that B₂₄, a new selfish variant showing accumulation, has completely replaced B₂ (a neutralized B chromosome incapable of accumulating) over the last few years in Torrox. In 1994, a new sample of individuals

revealed that the B₂₄ frequency had continued to increase, reaching its highest value ever recorded in a natural population of *E. plorans* (1.533) and that B₂ was no longer present. It seems, therefore, that the regeneration of polymorphism had already been completed in this population. If our theory regarding the dynamic evolution of the B-chromosome polymorphism is correct, B₂₄ should be neutralized within the next few years. In fact, the first evidence of suppressor genes against B₂₄ drive has already appeared: a small proportion of the crosses made by Zurita *et al.* (1998) showed a B₂₄ transmission rate close to a Mendelian one and they even found one female with one B chromosome to transmit this chromosome to only 15.2% of her offspring.

REFERENCES

- Adam, R. D. 1992 Chromosome-size variation in *Giardia lamblia*: the role of rDNA repeats. *Nucl. Acids Res.* **20**, 3057–3061.
- Alexander, M. 1981 Why microbial predators and parasites do not eliminate their prey and hosts. *A. Rev. Microbiol.* **35**, 113–133.
- Alfenito, M. R. & Birchler, J. A. 1993 Molecular characterization of a maize B-chromosome centric sequence. *Genetics* **135**, 589–597.
- Amos, A. & Dover, G. 1981 The distribution of repetitive DNAs between regular and supernumerary chromosomes in species of *Glossina* (tsetse): a two-step process in the origin of supernumeraries. *Chromosoma* **81**, 673–690.
- Anderson, R. M. & May, R. M. 1982 Coevolution of hosts and parasites. *Parasitology* **85**, 411–426.
- Battaglia, E. 1964 Cytogenetics of B chromosomes. *Caryologia* **17**, 245–299.
- Benazzi Lentati, G. 1970 Gametogenesis and egg fertilization in planarians. *Int. Rev. Cytol.* **27**, 101–179.
- Beukeboom, L. W. 1994a Bewildering Bs: an impression of the 1st B-chromosome conference. *Heredity* **73**, 328–336.
- Beukeboom, L. W. 1994b Phenotypic fitness effects of the selfish B chromosome, paternal sex ratio (PSR) in the parasitic wasp *Nasonia vitripennis*. *Evol. Ecol.* **8**, 1–24.
- Beukeboom, L. W. & Vrijenhoek, R. C. 1998 Evolutionary genetics and ecology of sperm-dependent parthenogenesis. *J. Evol. Biol.* **11**, 755–782.
- Beukeboom, L. W. & Werren, J. H. 1992 Population genetics of a parasitic chromosome: experimental analysis of PSR in subdivided populations. *Evolution* **46**, 1257–1268.
- Beukeboom, L. W., Seif, M., Mettenmeyer, T., Plowman, A. B. & Michiels, N. K. 1996 Paternal inheritance of B chromosomes in a parthenogenetic hermaphrodite. *Heredity* **77**, 646–654.
- Bigot, Y., Hamelin, M.-H. & Periquet, G. 1990 Heterochromatin condensation and evolution of unique satellite-DNA families in two parasitic wasp species: *Diadromus pulchellus* and *Eupelmus vuilleti* (Hymenoptera). *Mol. Biol. Evol.* **7**, 351–364.
- Blunden, R., Wilkes, T. J., Forster, J. W., Jiménez, M. M., Sandery, M. J., Karp, A. & Jones, R. N. 1993 Identification of the E3900 family, a 2nd family of rye chromosome-B specific repeated sequences. *Genome* **36**, 706–711.
- Bosemark, N. O. 1954 On accessory chromosomes in *Festuca pratensis*. II. Inheritance of the standard type of accessory chromosome. *Hereditas* **40**, 425–437.
- Bosemark, N. O. 1957 Further studies on accessory chromosomes in grasses. *Hereditas* **43**, 236–297.
- Bougourd, S. M. & Parker, J. S. 1979 The B chromosome system of *Allium schoenoprasum*. II. Stability, inheritance and phenotypic effects. *Chromosoma* **75**, 369–383.

- Bougourd, S. M., Plowman, A. B., Ponsford, N. R., Elias, M. L., Holmes, D. S. & Taylor, S. 1995 The case for unselfish B-chromosomes: evidence from *Allium schoenoprasum*. In *Kew Chromosome Conference IV* (ed. P. E. Brandham & M. D. Bennet), pp. 21–34. Kew, UK: Royal Botanic Gardens.
- Brockhouse, C., Bas, J. A. B., Fereday, R. M. & Strauss, N. A. 1989 Supernumerary chromosomes evolution in the *Simulium vernum* group (Diptera: Simuliidae). *Genome* **32**, 516–521.
- Bull, J. J., Molineux, I. J. & Rice, W. G. 1991 Selection of benevolence in a host–parasite system. *Evolution* **45**, 875–882.
- Butler, D. K. & Metznerberg, R. L. 1989 Premeiotic change of nucleolus organizer size in *Neurospora*. *Genetics* **122**, 783–791.
- Cabrero, J., Alché, J. D. & Camacho, J. P. M. 1987 Effects of B chromosomes of the grasshopper *Eyprepocnemis plorans* on nucleolar organizer regions activity. Activation of a latent NOR on a B chromosome fused to an autosome. *Genome* **29**, 116–121.
- Camacho, J. P. M., Cabrero, J., López-León, M. D. & Shaw, M. W. 1997a Evolution of a near-neutral B chromosome. In *Chromosomes today*, vol. 12 (ed. N. Henriques-Gil, J. S. Parker & M. J. Puertas), pp. 301–318. London: Chapman & Hall.
- Camacho, J. P. M., Shaw, M. W., López-León, M. D., Pardo, M. C. & Cabrero, J. 1997b Population dynamics of a selfish B chromosome neutralized by the standard genome in the grasshopper *Eyprepocnemis plorans*. *Am. Nat.* **149**, 1030–1050.
- Carlson, W. 1969 Factors affecting preferential fertilization in maize. *Genetics* **62**, 543–554.
- Carter, C. R. 1978 The cytology of *Brachycome*. 8. The inheritance, frequency and distribution of B chromosomes in *B. dichromosomatica* ($n = 2$), formerly in *B. lineariloba*. *Chromosoma* **67**, 109–121.
- Castro, A. J., Perfectti, F., Pardo, M. C., Cabrero, J., López-León, M. D. & Camacho, J. P. M. 1998 No harmful effects of a selfish B chromosome on several morphological and physiological traits in *Locusta migratoria* (Orthoptera, Acrididae). *Heredity* **80**, 753–759.
- Cebriá, A., Navarro, M. L. & Puertas, M. J. 1994 Genetic control of B chromosome transmission in *Aegilops speltoides* (Poaceae). *Am. J. Bot.* **81**, 1502–1507.
- Charlesworth, B. 1978 Model for evolution of Y chromosomes and dosage compensation. *Proc. Natl Acad. Sci. USA* **75**, 5618–5622.
- Charlesworth, B. & Lapid, A. 1989 A study of ten families of transposable elements on X chromosomes from a population of *Drosophila melanogaster*. *Genet. Res.* **54**, 113–126.
- Charlesworth, B., Langley, C. H. & Stephan, W. 1986 The evolution of restricted recombination and the accumulation of repeated DNA sequences. *Genetics* **112**, 947–962.
- Charlesworth, B., Lapid, A. & Canada, D. 1992a The distribution of transposable elements within and between chromosomes in a population of *Drosophila melanogaster*. I. Element frequencies and distribution. *Genet. Res.* **60**, 103–114.
- Charlesworth, B., Lapid, A. & Canada, D. 1992b The distribution of transposable elements within and between chromosomes in a population of *Drosophila melanogaster*. II. Inferences on the nature of selection against elements. *Genet. Res.* **60**, 115–130.
- Charlesworth, B., Sniegowski, P. & Stephan, W. 1994 The evolutionary dynamics of repetitive DNA in eukaryotes. *Nature* **371**, 215–220.
- Chen, X., Santhana Mariappan, S. V., Catasti, P., Ratliff, R., Moyzis, R. K., Laayoun, A., Smith, S. S., Morton Bradbury, E. & Gupta, G. 1995 Hairpins are formed by the single DNA strands of the fragile X triplet repeats: structure and biological implications. *Proc. Natl Acad. Sci. USA* **92**, 5199–5203.
- Clayton, D. H. & Tompkins, D. M. 1994 Ectoparasite virulence is linked to mode of transmission. *Proc. R. Soc. Lond. B* **256**, 211–217.
- Costanzi, C. & Pehrson, J. R. 1998 Histone macroH2A1 is concentrated in the inactive X chromosome of female mammals. *Nature* **393**, 599–601.
- Cuadrado, A. & Jouve, N. 1994 Highly repetitive sequences in B-chromosomes of *Secale cereale* revealed by fluorescence *in situ* hybridization. *Genome* **37**, 709–712.
- Dai, K., Gillies, C. B. & Dollin, A. E. 1994 Synaptonemal complex analysis of domestic sheep (*Ovis aries*) with Robertsonian translocations. III. Deficient pairing and NOR role in Massey III heterozygotes. *Genome* **37**, 802–808.
- Dawley, R. M. 1989 An introduction to unisexual vertebrates. In *Evolution and ecology of unisexual vertebrates* (ed. R. M. Dawley & J. P. Bogart). New York: University of the State of New York.
- Dherawattana, A. & Sadanaga, K. 1973 Cytogenetics of a crown rust-resistant hexaploid oat with 42 + 2 fragment chromosomes. *Crop Sci.* **13**, 591–594.
- Donald, T. M., Houben, A., Leach, C. R. & Timmis, J. N. 1997 Ribosomal RNA genes specific to the B chromosomes in *Brachycome dichromosomatica* are not transcribed in leaf tissue. *Genome* **40**, 674–681.
- Dover, G. A. & Riley, R. 1972 The prevention of pairing of homoeologous meiotic chromosomes of wheat by a genetic activity of supernumerary chromosomes of *Aegilops*. *Nature* **240**, 159–161.
- Eickbush, D. G., Eickbush, T. H. & Werren, J. H. 1992 Molecular characterization of repetitive DNA sequences from a B chromosome. *Chromosoma* **101**, 575–583.
- Epstein, H., James, T. C. & Singh, P. B. 1992 Cloning and expression of *Drosophila* HPI homologs from a mealybug *Planococcus citri*. *J. Cell. Sci.* **101**, 463–474.
- Ewald, P. 1987 Transmission modes and evolution of parasitism–mutualism continuum. *Ann. NY Acad. Sci.* **503**, 295–306.
- Ewald, P. W. & Schubert, J. 1989 Vertical and vector-borne transmission of insect endocytobiosis and the evolution of benignity. In *Insect endocytobiosis: morphology, physiology, genetics, evolution* (ed. W. Scwemmler & G. Gassner), pp. 21–35. Boca Raton, FL: CRC Press, Inc.
- Fox, D. P., Hewitt, G. M. & Hall, D. J. 1974 DNA replication and RNA transcription of euchromatic and heterochromatic chromosome regions during grasshopper meiosis. *Chromosoma* **45**, 43–62.
- Franks, T. K., Houben, A., Leach, C. R. & Timmis, J. N. 1996 The molecular organization of a B chromosome tandem repeat sequence from *Brachycome dichromosomatica*. *Chromosoma* **105**, 223–230.
- Fröst, S. 1958 Studies of the genetical effects of accessory chromosomes in *Centaurea scabiosa*. *Hereditas* **44**, 112–122.
- Fröst, S. 1969 The inheritance of accessory chromosomes in plants, especially in *Ranunculus acris* and *Phleum nodosum*. *Hereditas* **61**, 317–326.
- Garrido, M. A., Jamilena, M., Lozano, R., Ruiz Rejón, C., Ruiz Rejón, M. & Parker, J. S. 1994 RDNA site number polymorphism and NOR inactivation in natural populations of *Allium schoenoprasum*. *Genetica* **94**, 67–71.
- Garrido-Ramos, M. A., Jamilena, M., Lozano, R., Cárdenas, S., Ruiz Rejón, C. & Ruiz Rejón, M. 1995 Cytogenetic analysis of gilthead seabream *Sparus aurata* (Pisces, Perciformes), a deletion affecting the NOR in a hatchery stock. *Cytogenet. Cell Genet.* **68**, 3–7.
- Geiser, D. M., Arnold, M. L. & Timberlake, W. E. 1996 Wild chromosomal variants in *Aspergillus nidulans*. *Curr. Genet.* **29**, 293–300.
- Gilson, E., Perrin, D., Clement, J.-M., Szmelcman, S., Dassa, E. & Hofnung, M. 1986 Palindromic units from *E. coli* as binding sites for a chromoid-associated protein. *Fed. Eur. Biochem. Soc.* **206**, 323–328.
- González-Fernández, A., Navarrete, M. H., & De la Torre, C. 1993 Role for early replicating DNA in preventing nucleolus genesis in proliferating plant cells. *Protoplasma* **175**, 138–143.

- Green, D. M. 1988 Cytogenetics of the endemic New Zealand frog, *Leiopelma hochstetteri*: extraordinary supernumerary chromosome variation and a unique sex-chromosome system. *Chromosoma* **97**, 55–70.
- Green, D. M. 1990 Muller's ratchet and the evolution of supernumerary chromosomes. *Genome* **33**, 818–824.
- Green, D. M., Zeyl, C. W. & Sharbel, T. F. 1993 The evolution of hypervariable sex and supernumerary (B) chromosomes in the relict New Zealand frog, *Leiopelma hochstetteri*. *J. Evol. Biol.* **6**, 417–441.
- Gutknecht, J., Sperlich, D. & Bachmann, L. 1995 A species specific satellite DNA family of *Drosophila subsilvestris* appearing predominantly in B chromosomes. *Chromosoma* **103**, 539–544.
- Guttman, D. S. & Charlesworth, D. 1998 An X-linked gene with a degenerate Y-linked homologue in a dioecious plant. *Nature* **393**, 263–266.
- Hackstein, J. H. P., Hochstenbach, R., Hauschteck-Jungen, E. & Beukeboom, L. W. 1996 Is the Y chromosome of *Drosophila* an evolved supernumerary chromosome? *BioEssays* **18**, 317–323.
- Håkansson, A. 1954 Transmission of accessory chromosomes in *Poa alpina*. *Hereditas* **40**, 523–526.
- Henriques-Gil, N. 1984 El sistema de cromosomas accesorios de *Eyprepocnemis plorans* (Acrididae: Orthoptera). PhD thesis, Universidad Complutense de Madrid, Spain.
- Henriques-Gil, N. & Arana, P. 1990 Origin and substitution of B chromosomes in the grasshopper *Eyprepocnemis plorans*. *Evolution* **44**, 747–753.
- Henriques-Gil, N., Arana, P. & Santos, J. L. 1983 Spontaneous translocations between B chromosomes and the normal complement in the grasshopper *Eyprepocnemis plorans*. *Chromosoma* **88**, 145–148.
- Henriques-Gil, N., Santos, J. L. & Arana, P. 1984 Evolution of a complex polymorphism in the grasshopper *Eyprepocnemis plorans*. *Chromosoma* **89**, 290–293.
- Herre, E. A. 1993 Population structure and the evolution of virulence in nematode parasites of fig wasps. *Science* **259**, 1442–1445.
- Herrera, J. A., López-León, M. D., Cabrero, J., Shaw, M. W. & Camacho, J. P. M. 1996 Evidence for B chromosome drive suppression in the grasshopper *Eyprepocnemis plorans*. *Hereditas* **76**, 633–639.
- Hewitt, G. M. 1973a The integration of supernumerary chromosomes into the orthopteran genome. *Cold Spring Harb. Symp. Quant. Biol.* **38**, 183–194.
- Hewitt, G. M. 1973b Variable transmission rates of a B chromosome in *Myrmeleotettix maculatus* (Thunb.). *Chromosoma* **40**, 83–106.
- Hewitt, G. M. 1973c Evolution and maintenance of B chromosomes. *Chromosomes Today* **4**, 351–369.
- Hewitt, G. M. 1979 Grasshopper and crickets. In *Animal cytogenetics. 3. Insecta I Orthoptera* (ed. B. John), pp. 1–170. Berlin & Stuttgart, Germany: Gebrüder Borntraeger.
- Hiremath, S. C. & Murthy, H. N. 1986 The structure, stability and meiotic behavior of B chromosomes in *Guizotia scabra* (vis.) Chiov. Ssp. *Scabra* (Compositae). *Caryologia* **39**, 397–402.
- Hoepflich, P. D. 1977 Host-parasite relationships and the pathogenesis of infectious disease. In *Infectious diseases* (ed. P. D. Hoepflich), pp. 34–45. London: Harper & Row.
- Holliday, R. 1987 X-chromosome reactivation. *Nature* **327**, 661–662.
- Holmes, D. S. & Bougourd, S. M. 1989 B-chromosome selection in *Allium schoenoprasum*. II. Natural populations. *Hereditas* **63**, 83–87.
- Houben, A., Kynast, R. G., Heim, U., Hermann, H., Jones, R. N. & Forster, J. W. 1996 Molecular cytogenetic characterisation of the terminal heterochromatic segment of the B chromosome of rye (*Secale cereale*). *Chromosoma* **105**, 97–103.
- Houben, A., Leach, C. R., Verlin, D., Rofe, R. & Timmis, J. N. 1997a A repetitive DNA sequence common to the different B chromosomes of the genus *Brachycome*. *Chromosoma* **106**, 513–519.
- Houben, A., Belyaev, N. D., Leach, C. R. & Timmis, J. N. 1997b Differences in histone H4 acetylation and replication timing between A and B chromosomes of *Brachycome dichromosomatica*. *Chromosome Res.* **5**, 233–237.
- Ishak, B., Jaafar, H., Maetz, J. L. & Rumpler, Y. 1991 Absence of transcriptional activity of the B-chromosomes of *Apodemus peninsulae* during pachytene. *Chromosoma* **100**, 278–281.
- Jackson, R. C. & Newmark, K. P. 1960 Effects of supernumerary chromosomes on production of pigment in *Haploppappus gracilis*. *Science* **132**, 1316–1317.
- Jamilena, M., Ruíz-Rejón, C. & Ruíz-Rejón, M. 1994 A molecular analysis of the origin of the *Crepis capillaris* B chromosome. *J. Cell. Sci.* **107**, 703–708.
- Jamilena, M., Garrido-Ramos, M., Ruíz-Rejón, M., Ruíz-Rejón, C. & Parker, J. S. 1995 Characterisation of repeated sequences from microdissected B chromosomes of *Crepis capillaris*. *Chromosoma* **104**, 113–120.
- Jiménez, M. M., Romera, F., Gallego, A. & Puertas, M. J. 1995 Genetic control of the rate of transmission of rye B chromosomes. II. 0B × 2B crosses. *Hereditas* **74**, 518–523.
- John, U. P., Leach, C. R. & Timmis, J. N. 1991 A sequence specific to B chromosomes of *Brachycome dichromosomatica*. *Genome* **34**, 739–744.
- Johnson, N. A. 1997 Selfish genetic elements: long-range dynamics predicted by non-equilibrium models. *Trends Ecol. Evol.* **12**, 376–378.
- Jones, G. H., Whitehorn, J. A. F. & Albin, S. M. 1989 Ultrastructure of meiotic pairing in B chromosomes of *Crepis capillaris*. I. One-B and two-B pollen mother cells. *Genome* **32**, 611–621.
- Jones, J. D. & Flavell, R. B. 1983 Chromosomal structure and arrangement of repeated DNA sequences in the telomeric heterochromatin of *Secale cereale* and its relatives. *Cold Spring Harb. Symp. Quant. Biol.* **47**, 1209–1213.
- Jones, R. N. 1985 Are B chromosomes selfish? In *The evolution of genome size* (ed. T. Cavalier-Smith), pp. 397–425. London: Wiley.
- Jones, R. N. 1991 B-chromosome drive. *Am. Nat.* **137**, 430–442.
- Jones, R. N. 1995 Tansley review no. 85: B chromosomes in plants. *New Phytol.* **131**, 411–434.
- Jones, R. N. & Puertas, M. J. 1993 The B-chromosomes of rye (*Secale cereale* L.). In *Frontiers in plant science research* (ed. K. K. Dhir & T. S. Sareen), pp. 81–112. Delhi: Bhagwati Enterprises.
- Jones, R. N. & Rees, H. 1982 *B chromosomes*. New York: Academic Press.
- Keyl, H. G. & Hägele, K. 1971 B chromosomes bei *Chironomus*. *Chromosoma* **35**, 403–417.
- Kimura, M. & Kayano, H. 1961 The maintenance of supernumerary chromosomes in wild populations of *Lilium callosum* by preferential segregation. *Genetics* **46**, 1699–1712.
- Leach, C. R., Donald, T. M., Franks, T. K., Spiniello, S. S., Hanrahan, C. F. & Timmis, J. N. 1995 Organisation and origin of a B chromosome centromeric sequence from *Brachycome dichromosomatica*. *Chromosoma* **103**, 708–714.
- Leclair, S., Ansan-Melayah, D., Rouxel, T. & Balesdent, M. 1996 Meiotic behaviour of the minichromosome in the phytopathogenic ascomycete *Leptosphaeria maculans*. *Curr. Genet.* **30**, 541–548.
- Lin, M. S., Zhang, A. & Fujimoto, A. 1995 Asynchronous DNA replication between 15q11.2q12 homologs: cytogenetic evidence for maternal imprinting and delayed replication. *Hum. Genet.* **96**, 572–576.
- Lipsitch, M., Nowak, M. A., Ebert, D. & May, R. M. 1995 The population dynamics of vertically and horizontally transmitted parasites. *Proc. R. Soc. Lond. B* **260**, 321–327.
- López-Fernández, C., Mezzanotte, R. & Gosálvez, J. 1992 Autosomal, sex and B chromosomes in *Eyprepocnemis plorans* (Orthoptera) viewed with restriction endonuclease *in situ* digestion. *Hereditas* **68**, 365–372.

- López-León, M. D., Cabrero, J., Camacho, J. P. M., Cano, M. I. & Santos, J. L. 1992a A widespread B chromosome polymorphism maintained without apparent drive. *Evolution* **46**, 529–539.
- López-León, M. D., Pardo, M. C., Cabrero, J. & Camacho, J. P. M. 1992b Random mating and absence of sexual selection for B chromosomes in two natural populations of the grasshopper *Eyprepocnemis plorans*. *Heredity* **69**, 558–561.
- López-León, M. D., Cabrero, J., Pardo, M. C., Viseras, E., Camacho, J. P. M. & Santos, J. L. 1993 Generating high variability of B chromosomes in the grasshopper *Eyprepocnemis plorans*. *Heredity* **71**, 352–362.
- López-León, M. D., Neves, N., Schwarzacher, T., Heslop-Harrison, T. S., Hewitt, G. M. & Camacho, J. P. M. 1994 Possible origin of a B chromosome deduced from its DNA composition using double FISH technique. *Chromosome Res.* **2**, 87–92.
- McAllister, B. F. 1995 Isolation and characterization of a retroelement from B chromosome (PSR) in the parasitic wasp *Nasonia vitripennis*. *Insect Mol. Biol.* **4**, 253–262.
- McAllister, B. F. & Werren, J. H. 1997 Hybrid origin of a B chromosome (PSR) in the parasitic wasp *Nasonia vitripennis*. *Chromosoma* **106**, 243–253.
- McIntyre, C. L., Pereira, S., Moran, L. B. & Appels, R. 1990 New *Secale cereale* (rye) DNA derivatives for the detection of rye chromosome segments in wheat. *Genome* **33**, 635–640.
- McQuade, L. R., Hill, R. J. & Francis, D. 1994 B-chromosome systems in the greater glider, *Petauroides volans* (Marsupialia, Pseudocheiridae). 2. Investigation of B-chromosome DNA sequences isolated by micromanipulation and PCR. *Cytogenet. Cell Genet.* **66**, 155–161.
- McVean, G. T. 1995 Fractious chromosomes—hybrid disruption and the origin of selfish genetic elements. *BioEssays* **17**, 579–582.
- Marshall Graves, J. A. 1995 The origin and function of the mammalian Y chromosome and Y-borne genes—and evolving understanding. *BioEssays* **17**, 311–321.
- Martín-Alganza, A., Cabrero, J., López-León, M. D., Perfectti, F. & Camacho, J. P. M. 1997 Supernumerary heterochromatin does not affect several morphological and physiological traits in the grasshopper *Eyprepocnemis plorans*. *Heredity* **126**, 187–189.
- Matzke, M., Varga, F., Berger, H., Scherthaner, J., Schweizer, D., Mayr, B. & Matzke, A. J. M. 1990 A 41–42 tandemly repeated sequence isolated from nuclear envelopes of chicken erythrocytes is located predominantly on microchromosomes. *Chromosoma* **99**, 131–137.
- May, R. M. & Anderson, R. M. 1983 Epidemiology and genetics in the coevolution of parasites and hosts. *Proc. R. Soc. Lond. B* **219**, 2381–2413.
- Miao, V. P., Covert, S. F. & VanEtten, H. D. 1991a A fungal gene for antibiotic resistance on a dispensable ('B') chromosome. *Science* **254**, 1773–1776.
- Miao, V. P., Matthews, D. E. & VanEtten, H. D. 1991b Identification and chromosomal locations of a family of cytochrome P-450 genes for pisatin detoxification in the fungus *Nectria haematococca*. *Mol. Gen. Genet.* **226**, 214–223.
- Mills, D. & McCluskey, K. 1990 Electrophoretic karyotypes of fungi: the new cytology. *Mol. Plant–Microbe Int.* **3**, 351–357.
- Mitas, M., Yu, A., Dill, J., Kamp, T. J., Chambers, E. J. & Haworth, I. S. 1995 Hairpin properties of single-stranded DNA containing a GC-rich triplet repeat: (CTG)₁₅. *Nucl. Acids Res.* **23**, 1050–1059.
- Mitchell McGrath, J. & Helgeson, J. P. 1998 Differential behavior of *Solanum brevidens* ribosomal DNA loci in a somatic hybrid and its progeny with potato. *Genome* **41**, 435–439.
- Montgomery, E. A., Huang, S.-M., Langley, C. H. & Judd, B. H. 1991 Chromosome rearrangement by ectopic recombination in *Drosophila melanogaster*: genome structure and evolution. *Genetics* **129**, 1085–1098.
- Müntzing, A. 1954 Cytogenetics of accessory chromosomes (B-chromosomes). *Caryologia* **56**, 282–301.
- Nanda, I., Feichtinger, W., Schmid, M., Schröder, J. H., Zischler, H. & Epplen, J. T. 1990 Simple repetitive sequences are associated with differentiation of the sex chromosomes in the guppy fish. *J. Mol. Evol.* **30**, 456–462.
- Nanda, I., Scharl, M., Feichtinger, W., Epplen, J. T. & Schmid, M. 1992 Early stages of sex chromosome differentiation in fish as analysed by simple repetitive DNA sequences. *Chromosoma* **101**, 301–310.
- Nanda, I., Scharl, M., Epplen, J. T., Feichtinger, W. & Schmid, M. 1993 Primitive sex chromosomes in poeciliid fishes harbor simple repetitive DNA sequences. *J. Exp. Zool.* **265**, 301–308.
- Neves, N., Barão, A., Castilho, A., Silva, M., Morais, L., Carvalho, V., Viegas, W. & Jones, R. N. 1992 Influence of DNA methylation on rye B-chromosome nondisjunction. *Genome* **35**, 650–652.
- Nur, U. 1962 A supernumerary chromosome with an accumulation mechanism in the lecanoid genetic system. *Chromosoma* **13**, 249–271.
- Nur, U. 1966 Harmful B chromosomes in a mealy bug population. *Genetics* **54**, 1225–1238.
- Nur, U. 1969 Mitotic instability leading to an accumulation of B-chromosomes in grasshoppers. *Chromosoma* **27**, 1–19.
- Nur, U. 1977 Maintenance of a 'parasitic' B chromosome in the grasshopper *Melanoplus femur-rubrum*. *Genetics* **87**, 499–512.
- Nur, U. 1978 Asymmetrically heteropycnotic X chromosomes in the grasshopper *Melanoplus femur-rubrum*. *Chromosoma* **68**, 165–185.
- Nur, U. & Brett, B. L. H. 1985 Genotypes suppressing meiotic drive of a B chromosome in the mealy bug *Pseudococcus obscurus*. *Genetics* **110**, 73–92.
- Nur, U. & Brett, B. L. H. 1987 Control of meiotic drive of B chromosomes in the mealy bug *Pseudococcus affinis* (*obscurus*). *Genetics* **115**, 499–510.
- Nur, U. & Brett, B. L. H. 1988 Genotypes affecting the condensation and transmission of heterochromatic B chromosomes in the mealy bug *Pseudococcus affinis*. *Chromosoma* **96**, 205–212.
- Oliver, J. L., Posse, F., Martínez-Zapater, J. M., Enríquez, A. M. & Ruíz Rejón, M. 1982 B chromosomes and E1 isoenzyme activity in mosaic bulbs of *Scilla autumnalis*. *Chromosoma* **85**, 399–403.
- Östergren, G. 1945 Parasitic nature of extra fragment chromosomes. *Bot. Notiser* **2**, 157–163.
- Pardo, M. C., López-León, M. D., Cabrero, J. & Camacho, J. P. M. 1994 Transmission analysis of mitotically unstable B chromosomes in *Locusta migratoria*. *Genome* **37**, 1027–1034.
- Parker, J. S., Taylor, S. & Ainsworth, C. C. 1982 The B chromosome system of *Hypochoeris maculata*. III. Variation in B-chromosome transmission rates. *Chromosoma* **85**, 229–310.
- Peacock, W. J., Dennis, E. J., Rhoades, M. M. & Pryor, A. 1981 Highly repeated DNA sequence limited to knob heterochromatin in maize. *Proc. Natl Acad. Sci. USA* **78**, 4490–4494.
- Peppers, J. A., Wiggins, L. E. & Baker, R. J. 1997 Nature of B chromosomes in the harvest mouse *Reithrodontomys megalotis* by fluorescence *in situ* hybridization (FISH). *Chromosome Res.* **5**, 475–479.
- Peters, G. B. 1981 Germ line polysomy in the grasshopper *Atractomorpha similis*. *Chromosoma* **81**, 593–617.
- Pigozzi, M. I. & Solari, A. J. 1998 Germ cell restriction and regular transmission of an accessory chromosome that mimics a sex body in the zebra finch, *Taeniopygia guttata*. *Chromosome Res.* **6**, 105–113.
- Plowman, A. B. & Bougourd, S. M. 1994 Selectively advantageous effects of B chromosomes on germination behavior in *Allium schoenoprasum* L. *Heredity* **72**, 587–593.

- Puertas, M. J., González-Sánchez, M., Manzanero, S., Romera, F. & Jiménez, M. M. 1998 Genetic control of the rate of transmission of rye B chromosomes. IV. Localization of the genes controlling B transmission rate. *Heredity* **80**, 209–213.
- Pukkila, P. J. & Skrzynia, C. 1993 Frequent changes in the number of reiterated ribosomal RNA genes throughout the life cycle of the basidiomycete *Coprinus cinereus*. *Genetics* **133**, 203–211.
- Reed, K. M., Beukeboom, L. W., Eickbush, D. G. & Werren, J. H. 1994 Junctions between repetitive DNAs on the PSR chromosome of *Nasonia vitripennis*: association of palindromes with recombination. *J. Mol. Evol.* **38**, 352–362.
- Romera, F., Jiménez, M. M. & Puertas, M. J. 1991 Genetic control of the rate of transmission of rye B chromosomes. I. Effects in 2B × 0B crosses. *Heredity* **66**, 61–65.
- Rosato, M., Chiavarino, A. M., Naranjo, C. A., Puertas, M. J. & Poggio, L. 1996 Genetic control of B chromosome transmission rate in *Zea mays* ssp. *Mays* (Poaceae). *Am. J. Bot.* **83**, 1107–1112.
- Ruiz-Rejón, M., Posse, F. & Oliver, J. L. 1980 The B chromosome system of *Scilla autumnalis* (Liliaceae): effects at the isozyme level. *Chromosoma* **79**, 341–348.
- Ruiz-Rejón, M., Ruíz-Rejón, C. & Oliver, J. L. 1987 La evolución de los cromosomas B: existen cromosomas egoístas? *Investigación y Ciencia* **133**, 92–101.
- Ryan, S. L., Saul II, G. B. & Conner, G. W. 1985 Aberrant segregation of R-locus genes in male progeny from incompatible crosses in *Mormoniella*. *J. Hered.* **76**, 21–26.
- Ryan, S. L., Saul II, G. B. & Conner, G. W. 1987 Separation of factors containing R-locus genes in *Mormoniella* stocks derived from aberrant segregation following incompatible crosses. *J. Hered.* **78**, 273–275.
- Sandery, M. J., Forster, J. W., Blunden, R. & Jones, R. N. 1990 Identification of a family of repeated sequences on the rye B chromosome. *Genome* **33**, 908–913.
- Sapre, A. B. & Deshpande, D. S. 1987 Origin of B chromosomes in *Coix* L. through spontaneous interspecific hybridization. *J. Hered.* **78**, 191–196.
- Schartl, M., Nanda, I., Schlupp, I., Wilde, B., Epplen, J. T., Schmidt, M. & Parzefall, J. 1995 Incorporation of subgenomic amounts of DNA as compensation for mutational load in a gynogenetic fish. *Nature* **373**, 68–71.
- Sharbel, T. F., Pijnacker, L. P. & Beukeboom, L. W. 1997 Multiple supernumerary chromosomes in the pseudogamous parthenogenetic flatworm, *Polycelis nigra*: lineage markers or remnants of genetic leakage? *Genome* **40**, 850–856.
- Sharbel, T. F., Green, D. M. & Houben, A. 1998 B chromosome origin in the endemic New Zealand frog *Leiopelma hochstetteri* through sex chromosome devolution. *Genome* **41**, 14–22.
- Shaw, M. W. 1984 The population genetics of the B-chromosome polymorphism of *Myrmeleotettix maculatus* (Orthoptera: Acrididae). *Biol. J. Linn. Soc.* **23**, 77–100.
- Shaw, M. W. & Hewitt, G. M. 1985 The genetic control of meiotic drive acting on the B chromosome of *Myrmeleotettix maculatus* (Orthoptera: Acrididae). *Heredity* **54**, 259–268.
- Shaw, M. W. & Hewitt, G. M. 1990 B chromosomes, selfish DNA and theoretical models: where next? In *Oxford surveys in evolutionary biology*, vol. 7 (ed. D. Futuyma & J. Antonovics), pp. 197–223. Oxford University Press.
- Shaw, M. W., Hewitt, G. M. & Anderson, D. A. 1985 Polymorphism in the rates of meiotic drive acting on the chromosome of *Myrmeleotettix maculatus*. *Heredity* **55**, 61–68.
- Stark, E. A., Connerton, I., Bennet, S. T., Barnes, S. R., Parker, J. S. & Forster, J. W. 1996 Molecular analysis of the structure of the maize B-chromosome. *Chromosome Res.* **4**, 15–23.
- Staub, R. W. 1987 Leaf striping correlated with the presence of B chromosomes in maize. *J. Hered.* **78**, 71–74.
- Steinemann, M. & Steinemann, S. 1991 Preferential Y chromosomal location of TRIM, a novel transposable element of *Drosophila miranda*, *obscura* group. *Chromosoma* **101**, 169–179.
- Steinemann, M. & Steinemann, S. 1992 Degenerating Y chromosome of *Drosophila miranda*: a trap for retrotransposons. *Proc. Natl Acad. Sci. USA* **89**, 7591–7595.
- Steinemann, M. & Steinemann, S. 1997 The enigma of Y chromosome degeneration: TRAM, a novel retrotransposon is preferentially located on the neo-Y chromosome of *Drosophila miranda*. *Genetics* **145**, 261–266.
- Steinemann, M., Steinemann, S. & Lottspeich, F. 1993 How Y chromosomes become genetically inert. *Proc. Natl Acad. Sci. USA* **90**, 5737–5741.
- Stephan, W. 1987 Quantitative variation and chromosomal location of satellite DNAs. *Genet. Res.* **50**, 41–52.
- Talavera, M., López-León, M. D., Cabrero, J. & Camacho, J. P. M. 1990 Male germ line polysomy in the grasshopper *Chorthippus binotatus*: extra chromosomes are not transmitted. *Genome* **33**, 384–388.
- Thomas, J. H. 1995 Genomic imprinting proposed as a surveillance mechanism for chromosome loss. *Proc. Natl Acad. Sci. USA* **92**, 480–482.
- Tzeng, T. H., Lyngholm, L. K., Ford, C. F. & Bronson, C. R. 1992 A restriction fragment length polymorphism and electrophoretic karyotype of the fungal maize pathogen *Cochliobolus heterostrophus*. *Genetics* **130**, 81–96.
- Viotti, A., Privitera, E., Sala, E. & Pogna, N. 1985 Distribution and clustering of two highly repeated sequences in the A and B chromosomes of maize. *Theor. Appl. Genet.* **70**, 234–239.
- Vogel, J. M., Nieto, M. C., Fishcher, A. & Goodenow, R. S. 1990 Overlapping palindromic sequences associated with somatic deletion and meiotic recombination of MHC class I genes. *Mol. Immunol.* **27**, 875–886.
- Werren, J. H. 1991 The paternal-sex-ratio chromosome of *Nasonia*. *Am. Nat.* **137**, 392–402.
- Werren, J. H. & Beukeboom, L. W. 1993 Population genetics of a parasitic chromosome: theoretical analysis of PSR in subdivided populations. *Am. Nat.* **142**, 224–241.
- Werren, J. H., Nur, U. & Eickbush, D. G. 1987 An extrachromosomal factor causing loss of paternal chromosomes. *Nature* **327**, 75–76.
- White, M. J. D. 1973 *Animal cytology and evolution*, 3rd edn. London: Cambridge University Press.
- Wilkes, T. M., Francki, M. G., Langidge, P., Karp, A., Jones, R. N. & Forster, J. W. 1995 Analysis of rye B-chromosome structure using fluorescence *in situ* hybridization (FISH). *Chromosome Res.* **3**, 466–472.
- Wilson, E. B. 1907 The supernumerary chromosomes of Hemiptera. *Science* **26**, 870–871.
- Zeyl, C. W. & Bell, G. 1996 Symbiotic DNA in eukaryotic genomes. *Trends Ecol. Evol.* **11**, 10–14.
- Zeyl, C. W. & Green, D. M. 1992 Heteromorphism for a highly repeated sequence in the New Zealand frog *Leiopelma hochstetteri*. *Evolution* **46**, 1891–1899.
- Zurita, S., Cabrero, J., López-León, M. D. & Camacho, J. P. M. 1998 Polymorphism regeneration for a neutralized selfish B chromosome. *Evolution* **52**, 274–277.